

Can Cosmic Rays Heat the Intergalactic Medium?

Saumyadip Samui, Kandaswamy Subramanian, Raghunathan Srianand
*IUCAA, Post Bag 4, Ganeshkhind, Pune - 411 007, India**
 (Dated:)

Supernova explosions in the early star forming galaxies will accelerate cosmic rays (CRs). CRs are typically confined in the collapsed objects for a short period before escaping into the intergalactic medium (IGM). Galactic outflows can facilitate this escape by advecting CRs into the IGM. An outflow that results in a termination shock can also generate more CRs. We show that the CR protons from the above processes can significantly affect the thermal history of the IGM. Within plausible range of parameters, cosmic ray heating can compensate for adiabatic cooling and explain the measured IGM temperature at redshifts $z \sim 2 - 4$, even with early reionization.

PACS numbers: 98.62.Ra, 98.70.Sa, 96.40.-z, 98.62.-g, 98.80.-k

It is widely believed that the temperature of the low density intergalactic medium retains some memory of the reionization process [1]. Analysis of Ly α absorption spectra in the framework of hydrodynamical simulations and semi-analytic models [2] suggest that, the mean temperature of the IGM is in the range $10^4 \leq T(\text{K}) \leq 3 \times 10^4$ for $2 \leq z \leq 4$. This has been used to argue that hydrogen reionization occurred below $z \simeq 9$ [3]. Had the reionization occurred at higher z , adiabatic cooling would have brought the temperature much below the measured value at $z = 4$. The detection of a strong Gunn-Peterson trough and the sizes of ionized regions around the highest z QSOs [4], are consistent with a large neutral hydrogen fraction at $z \simeq 6$. The CMB data from the WMAP satellite [5], however, points to an earlier epoch of reionization in the range $11 < z < 30$. Thus either new heating sources (other than photo-heating) or multiple episodes of reionization with heating at low z provided by, say, the He II reionization induced by QSOs [6], are needed to explain the IGM temperature at $z \sim 2 - 4$. Here, we consider the effect of cosmic ray protons on the thermal history of the IGM.

Cosmic rays are expected to be generated and accelerated from the shocks created by the exploding supernovae (SNe) in the star forming regions, termination shocks created by outflows from galaxies and accretion shocks during structure formation. The properties of CRs in our Galaxy is well documented [7, 8, 9, 10]. The energy density in the proton component is about 1 eV cm^{-3} , and CRs are thought to be confined to the Galactic disk for about 10^7 yr before escaping, presumably into the IGM. It is widely accepted that $\geq 15\%$ of the average SNe energy must go into accelerating the CRs so that the flux density of CRs can be maintained at the observed value [10]. Low energy CR protons play an important role in the ionization and thermal state of the gas in the interstellar medium (ISM) of normal galaxies. The possibility that they could also play an important role in determining the thermal history of the IGM has been considered sporadically in the literature [11, 12]. The existence of good observational constraints on (i) the global star for-

mation rate (SFR) density as a function of z (ii) epoch of reionization and (iii) temperature of the IGM makes it appropriate to re-examine this issue using a model that self consistently computes the generation and spectral evolution of CRs in the IGM. We do this here in the framework of a standard flat Λ CDM model ($\Omega_m = 0.27$, $\Omega_b = 0.04$ and $H_0 = 70 \text{ km s}^{-1} \text{ Mpc}^{-1}$).

Consider CR protons produced from the SNe explosions. To compute the rate of energy injected into CR protons from early star forming galaxies, one needs to know the SFR and the number of SNe explosions, f_{SN} , per solar mass of forming stars. We use the observed global comoving SFR density [13], $R(z)$ measured in units of $M_\odot \text{ yr}^{-1} \text{ Mpc}^{-3}$ and approximated by

$$R(z) = 7.5 \times 10^{-4} \eta z^{2.5} e^{-0.8(z-3.5)}. \quad (1)$$

Here, η , in the range 5 to 9, is the unknown scaling factor used to correct the effect of dust reddening [14]. We shall adopt $\eta = 7$ in our models. Further, using the ‘‘Starburst99’’ code [15] we get $f_{SN} = 8 \times 10^{-3}$ for Salpeter IMF (with $M_{min} = 0.1 M_\odot$, $M_{max} = 100 M_\odot$) and $f_{SN} = 1.7 \times 10^{-2}$ for a top-hat mass function with $M_{min} = 40 M_\odot$, $M_{max} = 100 M_\odot$ (denoted as ‘‘top-heavy’’). We assume that the energy released from a single SNe is 10^{51} erg and a fraction $\epsilon = 0.15$ of this is utilized for accelerating the CR protons. The average rate of energy injected in the CR protons is then,

$$\dot{E}_{CR}(z) = 10^{-30} \epsilon R f_{SN} (1+z)^3 \text{ erg s}^{-1} \text{ cm}^{-3}. \quad (2)$$

The cosmic ray proton spectrum from the SNe is in general taken to be a power law in momentum space, with $n(p) dp \propto p^{-q} dp$, with $q = 2.2$ as a typical value [7]. We can then compute the average rate of CRs generated in star forming regions as

$$\frac{dn(p, z)}{dt} dp = \dot{N}_0 p^{-q} dp \quad (3)$$

where the normalization $\dot{N}_0$ for a given z is fixed by integrating $E(p) \dot{N}_0 p^{-q} dp$ using a lower energy cut-off [7] of 10 keV, and equating the result to $\dot{E}_{CR}(z)$. Here $E(p)$




FIG. 1: Cosmic rays spectrum originating from SNe (dotted line) and the modified spectra (solid lines) escaping from galaxy. The curves A and B are for $x = 10$ and 3 g cm^{-2} respectively. The curve C is for the advection model, where the spectrum A has been modified by decreasing the momentum of each CR particle by a factor 10. Curve D is the CR spectrum originating from the termination shock with a sharp cut-off at 3 MeV. For illustrative purpose the injected $Q(E, z)$ is normalized using R at $z = 15$.

is the kinetic energy of a CR proton corresponding to a momentum p . This injected spectrum will be modified as the CRs propagate first within the high redshift star forming galaxy and then into the IGM. The evolution of the velocity $v = \beta c$ of the CR particle due to collisions with free electrons, hydrogen ionization, and the adiabatic expansion of the universe is given by [7, 12],

$$\begin{aligned}
 -\frac{d\beta}{dt} = & 3.27 \times 10^{-16} n_e(z) \frac{(1-\beta^2)^{3/2}}{\beta} \frac{\beta^2}{x_m^3 + \beta^3} \\
 & + 1.94 \times 10^{-16} n_{HI}(z) \frac{(1-\beta^2)^{3/2}}{\beta} \\
 & \times \frac{2\beta^2}{[(0.01)^3 + 2\beta^3]} [1 + 0.0185 \ln \beta \Theta(\beta - 0.01)] \\
 & - \frac{\beta(1-\beta^2)}{1+z} \frac{dz}{dt}.
 \end{aligned} \quad (4)$$

Here, n_e and n_{HI} are the densities of electrons and neutral hydrogen respectively in units of cm^{-3} . Θ is Heaviside function, $x_m = 0.0286 (T/2 \times 10^6 K)^{1/2}$ and T is the temperature of the gas. The energy losses are dominated by low energy CRs for $\beta > x_m$. Typically a CR particle of energy E will lose all its energy to the electrons in a time, $t_L \simeq 1.5 \times 10^8 (E/3\text{MeV})^{1.5} (10^{-4}/n_e)$ yr, after traveling a distance $d_L \sim 4(E/3\text{MeV})^2 (10^{-4}/n_e)$ Mpc.

The escape of the CRs from the star forming galaxies will depend on the structure and strength of its magnetic field, the gas density (ρ) and outflows etc. We consider three possible scenarios for the emerging CR spectrum: (I) CRs are confined for a period t_c to the galaxy and then escape into the IGM, (II) after a confinement time t_c ,

CRs are advected by galactic outflows into the IGM, and (III) CRs are generated in termination shocks created by the outflows.

In case (I), we compute the emergent spectrum from the galaxy, taking into account the energy loss to electrons, as a function of grammage, $x \sim \rho c t_c \text{ g cm}^{-2}$, traversed by the CRs within the galaxy. For our Galaxy x is $\sim 10 \text{ g cm}^{-2}$ [9]. For the high z galaxies, x could be smaller, because the global star formation rate density is expected to be dominated by smaller mass objects that may also be ‘magnetically’ younger and will confine CRs for a smaller period. The injected spectrum as well as modified spectra (defined as the rate $Q(E, z)dE = (dn/dt)(dt/dz)dp$ per unit redshift) for $x = 10, 3 \text{ g cm}^{-2}$, are shown in the Fig. 1 and labeled as A and B. Larger and larger part of the low energy CR-spectrum is cut off due to energy losses as one increases x . As an example, when $x = 10 \text{ g cm}^{-2}$ protons of initial energy $\leq 160 \text{ MeV}$ will lose all their energy. Energy loss from protons with higher energy does lead to an universal low energy tail with $Q \propto E^{1/2}$. Since $dE/dt \propto E^{-1/2}$, the heating is still dominated by protons near the peak of Q .

High redshift Lyman break galaxies (LBGs) show signatures of strong outflows [16]. Such outflows, that are also required to explain the observed IGM metallicity, will help in advecting the CRs into the IGM. The adiabatic loss in such outflows can also lower the momentum (p) of the CR particles, with $p \propto \rho^{1/3}$, thereby shifting the spectrum to lower momentum and decreasing the total energy in CRs. Suppose a wind from a galaxy of 3 kpc size expands up to 30 kpc before CRs escape into the IGM. The average gas density will decrease by a factor 10^3 thereby reducing the momentum of CRs by a factor 10. The emergent spectrum for this case (our case (II)) (and $x = 10 \text{ g cm}^{-2}$) is also shown in Fig. 1 (see curve C).

In order for the CRs to influence the properties of the IGM, they should be released into the IGM within the Hubble time. This time-scale is confinement time for case (I) and advection time for case (II). In our Galaxy $t_c \sim 10^7$ yr in the disk. In the Galactic halo of say a radius r_H , the typical t_c is the diffusion time, $r_H^2/6D \sim 10^8$ yr, for $r_H \sim 10$ kpc and a diffusion coefficient $D = 5 \times 10^{28} \text{ cm}^2 \text{ s}^{-1}$ [8]. Thus for case (I) t_c is expected to be smaller than the Hubble time, even for the Galaxy. Similarly the advection time scale in an outflow of velocity typically seen in LBGs (i.e $\sim 100 \text{ km s}^{-1}$) to a radius 30 kpc as in our case (II) will be 3×10^8 yr, which is shorter than the Hubble time for $z \leq 15$.

In case (III) we consider the CRs produced from the termination shocks of the outflows. We assume that about 10% of the SNe energy is channeled into driving such an outflow, and a fraction ϵ of this goes into accelerating the CRs in the resulting termination shock. The spectrum of CRs emerging from the shock is again ex-

pected to be a power law in momentum space. As the wind termination shock is expected to form far out in the halo of the galaxy and propagate for most periods in the IGM itself, even low energy cosmic rays can be directly injected into the IGM. However the lowest energy CRs will lose their energy in the vicinity of the galaxy itself. For example, a 3 MeV CR proton will travel a proper distance $d_L \sim 10$ Mpc in the IGM at $z \sim 5$ before losing all its energy. While within a halo that has 200 times larger density, it will travel ~ 50 kpc. Thus we assume that the particles with energy $E \geq 3$ MeV will be able to escape from termination shocks and contribute to the heating of the IGM. The resulting spectrum with a sharp cut-off at 3 MeV is shown in Fig. 1 as case D.

The free propagation of CRs in the IGM will depend on the strength and coherence of the intergalactic magnetic field. For example in the case of ISM the streaming of CRs is limited by resonant scattering on self-generated Alfvén waves. It turns out that a coherent IGM magnetic field larger than $\sim 10^{-14}$ G is required for a streaming instability growth time-scale $< 10^8$ yr (Eq. 32 in [19]), so that there can be adequate growth of Alfvén waves, by say $z \sim 5$. Astrophysical battery mechanisms produce only tiny coherent IGM seed fields $\sim 10^{-23} - 10^{-19}$ G [17]. Early Universe generation mechanisms are not very compelling because the results are often exponentially sensitive to parameters involved [18]. Further, for $z \geq 2$, the self-confinement due to plasma waves generated by the streaming CRs [11] is inefficient because of the strong damping of waves by electron ion collisions. Here, we will assume that the CRs escaping the high z galaxies are not self confined, but can stream out freely in the IGM.

The spectrum of cosmic ray background at a given z is the sum of CR spectrum injected at z and that evolving from higher redshifts. The number density of CRs, $N_{CR}(\beta, z, z_0)d\beta$, in an interval $(\beta, \beta + d\beta)$, at any given epoch, z , is

$$N_{CR}(\beta, z, z_0) = \int_{z_0}^z dz_i \frac{dn(z_i, p_i)}{dz_i} \frac{dp_i}{d\beta_i} \frac{d\beta_i}{d\beta} \left(\frac{1+z}{1+z_i} \right)^3. \quad (5)$$

In order to compute N_{CR} one needs to know the β evolution and hence the ionization history of the IGM. We model this using a very simple reionization scenario with the aim to merely compute N_{CR} in a self-consistent setting rather than addressing the issues related to early reionization. The rate of change in the number density of ionized hydrogen (i.e. n_{HII}) can be written as,

$$\frac{dn_{HII}}{dt} = \dot{N}_\gamma + \int N_{CR}(\beta, z, z_0) cn_{HI} \sigma_{HI} \beta d\beta - \alpha_B n_{HII} n_e. \quad (6)$$

The right hand side of Eq. 6 gives the rates of photoionization, cosmic ray ionization and recombination respectively. We assume all the Lyman continuum photons




FIG. 2: Predicted temperature evolution of the IGM for different injected spectra given in Fig. 1. We follow the labeling convention as in Fig. 1. Temperature evolution for the adiabatic cooling is shown with a dotted curve. The data points are a compendium of observations presented in [2].

entering the IGM with a rate $\dot{N}_\gamma$ are used for ionizing H I. For a SFR of $1 M_\odot \text{ yr}^{-1}$ about 24000 s^{-1} and 4000 s^{-1} Lyman continuum photons are generated per baryon, for “top-heavy” and “Salpeter” IMF respectively. This is used with an escape fraction of 0.1 to compute $\dot{N}_\gamma$. σ_H is the cross-section for H I ionization by protons [20] and α_B is the case-B recombination coefficient. We find CR ionization is sub-dominant compared to the photoionization.

We assume “top-heavy” IMF prior to the epoch of reionization and “Salpeter” IMF after that. The value of n_{HII} at a given z is obtained using Eq. 6 and the analytical fit to the observed SFR density as a function of z (Eq. 1). We get the epoch of reionization (where $n_{HII} = n_H$) at $z_r \simeq 10$. Note that this is independent of our choice of η because while $R \propto \eta$ the UV escape fraction $f_e \propto 1/\eta$. However, as SNe rate increases with η the CR flux density will depend on η . Using the reionization model, and Eqs. 1 - 5 we compute the spectrum of CR at any given z , starting at $z_0 = 15$. Such spectra are used below to compute the heating of the IGM due to CRs.

In our model calculations we assume (i) IGM reaches an uniform temperature of $T = 3 \times 10^4$ K at $z = z_r$, (ii) heating by UV photons, Compton cooling by CMB and recombination cooling are negligible for $z < z_r$ and (iii) for simplicity the IGM is assumed to be of uniform density. Thus the thermal state of the IGM after reionization is described by,

$$\frac{d\mathcal{E}}{dz} = \frac{5\mathcal{E}}{(1+z)} - \Gamma_{CR} \quad (7)$$

with $\mathcal{E}$ is the thermal energy per unit volume of the IGM. The first term in the right hand side of Eq. 7 gives adiabatic cooling rate. The second term Γ_{CR} is the cosmic

rays heating rate given by

$$\begin{aligned}\Gamma_{CR}(z, z_0) &= \int \frac{dE(\beta)}{dz} N_{CR}(\beta, z, z_0) d\beta \\ &= \int \frac{dE(\beta)}{dt} \frac{dt}{dz} N_{CR}(\beta, z, z_0) d\beta.\end{aligned}\quad (8)$$

Here, dE/dt is obtained from the first term of Eq. 4. Using $\mathcal{E} = 3nkT/2$, where T is the temperature and n the number density of particles in the IGM, one can obtain

$$T(z) = T(z_r) \left(\frac{1+z}{1+z_r} \right)^2 \times \left[1 + \frac{(1+z_r)^2}{3n_0 k T(z_r)} \int_z^{z_r} \frac{\Gamma_{CR}(z', z_0)}{(1+z')^5} dz' \right], \quad (9)$$

where $n_0 = n_e(z=0)$.

The temperature evolution of the IGM for different injected spectra (shown in Fig. 1) computed using Eq. 9 are shown in Fig. 2. The temperature of the IGM goes well below the observed range due to adiabatic cooling in the absence of heating sources (dotted curve in Fig. 2). However for all the models discussed here, CR heating leads to, $T \geq 10^4 \text{K}$ in the redshift range $2 \leq z \leq 4$.

For case (I) the average grammage traversed by the CRs in the high z star forming galaxies has to be less than 10 g cm^{-2} seen in the Galaxy, in order to produce the observed mean IGM temperature (curves A and B in Fig. 2). A lower grammage is reasonable for the high z galaxies, as mentioned earlier. These models predict cosmic ray energy density $U_{cr} \sim 6 \times 10^{-3} \text{ eV cm}^{-3}$ at $z = 0$. Models with adiabatic losses (case (II)) or termination shocks (case (III)) produce higher temperatures, as they efficiently inject lower energy CRs into the IGM. At the same time they lead to overall lower cosmic ray energy densities of the order $1.5 \times 10^{-4} \text{ eV cm}^{-3}$ and $5.7 \times 10^{-4} \text{ eV cm}^{-3}$ for case (II) and (III) respectively at $z = 0$.

Models of shock acceleration [10] predict ϵ as high as 0.3 – 0.4. In such a case even a model with grammage as seen in our galaxy will produce higher temperatures than the observed mean. Simulations of pregalactic outflows [21], find up to 30% of the supernovae energy can go into winds. The resulting temperature in case (III) can thus be much higher. We considered CRs produced in the SNe and termination shocks, CRs generated in structure formation shocks [22] could provide additional heating. The major uncertainties are related to the extent to which the IGM is magnetized and confinement issues. More detailed computations in the framework of structure formation models are in progress.

We have explored the influence of CRs escaping from the early star forming regions on the thermal history of the IGM. The estimated IGM temperature evolution is sensitive to the star formation rate (η) and the confinement of CRs. Using a star formation rate density con-

strained by observations and plausible range of parameters, the observed temperature of the IGM at $2 \leq z \leq 4$ can be explained as due to the CR heating, even if the epoch of reionization is as early as $z = 10$. This implies that low energy CRs in the IGM can erase the thermal memory of reionization. In conclusion, cosmic rays do provide an important source for heating the intergalactic medium, that can explain the measured IGM temperatures at high redshifts.

SS thanks CSIR for providing support for this work. We thank A. Shukurov for helpful comments.

* Electronic address: samui@iucaa.ernet.in, kandu@iucaa.ernet.in, anar@iucaa.ernet.in

- [1] Miralda-Escude J. and Rees M. J., MNRAS **266**, 343 (1994); Hui L. and Gnedin N. Y., MNRAS **292**, 27 (1997); Haehnelt M. G. and Steinmetz, M., MNRAS **298**, L21 (1998).
- [2] Schaye J., et. al., MNRAS **318**, 817 (2000); Ricotti M., Gnedin N. Y. and Shull J. M., ApJ **534**, 41 (2000); Choudhury T., Srianand R. and Padmanabhan T., ApJ **559**, 29 (2001); McDonald P., et. al., ApJ **562**, 52 (2001); Zaldarriaga M., Hui L. and Tegmark M., ApJ **557**, 519(2001).
- [3] Theuns T., et. al., ApJ **567**, L103 (2002).
- [4] Fan X., et. al., AJ **123**, 1247 (2002); Wyithe J. S. B. and Loeb A., Nature **427**, 815 (2004).
- [5] Kogut A., et. al., ApJS **148**, 161 (2003).
- [6] Hui L. and Haiman, Z., ApJ **596**, 9 (2003).
- [7] Schlickeiser R., Cosmic Ray Astrophysics, Berlin, Springer-Verlag, (2000).
- [8] Berezhinskii, V. S., et. al., Astrophysics of Cosmic Rays, Amsterdam, North-Holland, (1990); Berezhinsky V. S., Blasi P., Ptuskin V. S., ApJ **487**, 529 (1997).
- [9] Dogiel V. A., Schonfelder, V. and Strong A. W., ApJ **572**, L157 (2002).
- [10] Hillas A. M., J. Phys. G: Nucl. Part. Phys. **31**, R95 (2005).
- [11] Ginzburg V. L. and Ozernoi L. M., Sov. Astr. **9**, 726 (1966).
- [12] Nath B. B. and Biermann P. L., MNRAS **265**, 241 (1993).
- [13] Bouwens R. J., et. al., astro-ph/0503116.
- [14] Adelberger K. L. and Steidel C. C., ApJ **544**, 218 (2000).
- [15] Leitherer C., et. al., ApJS **123**, 3 (1999).
- [16] Shapley A. E. et al., ApJ, **588**, 65 (2003).
- [17] Subramanian, K., Narasimha, D., Chitre, S. M., MNRAS, **271**, 15 (1994); Gnedin, N. Y., Ferrara, A., Zweibel, E. G., ApJ, **539**, 505 (2000).
- [18] Widrow L. M., Rev. Mod. Phys. **74**, 775 (2002).
- [19] Zweibel E. G., ApJ **587**, 625 (2003).
- [20] Spitzer L. Jr. and Tomasko M. G., ApJ **152**, 971 (1968).
- [21] Mori, M., Ferrara, A. and Madau, P., ApJ, **571**, 40 (2003).
- [22] Miniati, F., et al., ApJ, **559**, 59 (2001); Furlanetto R and Loeb A. ApJ, **611**, 642 (2004); Kang H., Jones T. W. ApJ, **620**, 44 (2005) and references therein.